

Case 3244

TERMOPSIDAE Holmgren, 1911, *Termopsis* Heer, 1849 and *Mioterмес* Rosen, 1913 (Insecta, Isoptera): proposed conservation of prevailing usage by the designation of *Termopsis breinii* Heer, 1849 as the type species of *Termopsis*

Michael S. Engel

*Division of Entomology, Natural History Museum, Snow Hall,
1460 Jayhawk Boulevard, University of Kansas, Lawrence,
Kansas 66045-7523, U.S.A.*

Kumar Krishna and Christopher Boyko

*Division of Invertebrate Zoology, American Museum of Natural History,
Central Park West at 79th Street, New York, N.Y. 10024-5192, U.S.A.*

Abstract. The purpose of this application, under Article 70.2 of the Code, is to conserve the current usage of the generic names *Termopsis* Heer, 1849 and *Mioterмес* Rosen, 1913 and the family name TERMOPSIDAE Holmgren, 1911 for well known groups of termites. Both nominal genera have the same type species, *Termopsis procerus* Heer, 1849, and it is proposed that this problem of synonymy be resolved by designation of *Termopsis breinii* Heer, 1849 as the type species of *Termopsis*.

Keywords. Nomenclature; taxonomy; Isoptera; TERMOPSIDAE; *Termopsis*; *Mioterмес*; *Termopsis breinii*; *Termopsis procerus*; termites.

1. Heer (1849, p. 23) named the termite genus *Termopsis* (at that time as a subgenus of *Termes* Linnaeus, 1758) and included five new fossil species, two of which were *Termopsis breinii* Heer, 1849 (p. 31) and *T. procerus* Heer, 1849 (p. 23), in Middle Eocene Baltic amber. Heer did not select any of the five species as the type species of *Termopsis*. Holmgren (1911, p. 35) established the family-group name TERMOPSIDAE based on *Termopsis*.

2. Handlirsch (1907, p. 698) designated *Termopsis procerus* Heer, 1849 as the type species of *Termopsis* and was the first author to explicitly designate a type species from one of Heer's originally included five species. Cockerell (1916, p. 138) also claimed to have designated *T. procerus* as the type of *Termopsis*.

3. Banks (in Banks & Snyder, 1920, p. 9) selected *Termopsis insignis* Heer, 1849 as the type species of *Termopsis* but, although he correctly chose one of Heer's original species, his action was not valid since Handlirsch (1907, p. 698) had already designated *T. procerus* as the type species.

4. Hagen (1854, p. 222) included only *Termopsis breinii* Heer in *Termopsis*, transferring four of Heer's original five species to the genus *Hodotermes* Hagen, 1853. Later, Hagen (in Pictet-Baraban & Hagen, 1856, p. 51; 1858a, p. 32 and 1858b, p. 12) again included only *T. breinii* of Heer's original five species in *Termopsis*. Although

Hagen never specifically mentioned *T. breinii* as the type species of *Termopsis*, his elimination of all species except *T. breinii* was interpreted later by Emerson (1933, p. 165) as the fixation of a type species. Article 69.4 states that 'elimination of all but one of the originally included nominal species from a nominal genus or subgenus does not in itself constitute type fixation'. Had Handlirsch (1907) not already designated *T. procerus* Heer, 1849 as the type species of *Termopsis* (and had Banks (1920) not chosen *T. insignis*), Emerson's (1933, p. 165) subsequent acceptance of *T. breinii* as the type species of *Termopsis* would have been valid under Article 69.1.1. This Article states that 'in the absence of a prior type fixation for a nominal genus or subgenus, an author is deemed to have designated one of the originally included nominal species as type species, if he or she states (for whatever reason, right or wrong) that it is the type or type species'.

5. Girard (1879, p. 270) selected *T. angusticollis* Hagen, 1858a as the type species of *Termopsis*. Similarly, Wasmann (1897, p. 149) selected *Termes occidentis* Walker, 1853 as the type species of *Termopsis*. Since neither *T. angusticollis* nor *T. occidentis* was originally included in *Termopsis* by Heer (1849), these designations are invalid under Article 67.2.

6. Rosen (1913, p. 325) described a new termite genus *Miotermes*, with *T. procerus* Heer, 1849 as the type species by original designation and monotypy and transferred the species to the family MASTOTERMITIDAE, a family proposed by Desneux (1904) for the primitive Australian termite *Mastotermes darwiniensis* Froggatt, 1897 (see Opinion 1808, 1995, BZN 52: 206).

7. However, Snyder (1949, p. 360), in a catalog of the termite species of the world, followed Emerson's (1933) conclusion concerning Hagen's various actions (1854, 1856, 1858a, 1858b) and listed *T. breinii* as the type species of *Termopsis*.

8. All authors since Snyder (1949) have followed the invalid interpretation of Emerson (1933) and have considered *Termopsis* to apply to *T. breinii* and related species. They have employed the family-group name TERMOPSIDAE Holmgren, 1911 (p. 35) for *Termopsis* (sensu current usage) and its relatives in an extensive systematic (e.g., Wilson, 1971; Thorne & Carpenter, 1992; Nel & Paicheler, 1993; Weitschat & Wichard, 1998, 2002; Krishna & Grimaldi, 2000; Thorne, Grimaldi, & Krishna, 2000), biological (Weidner, 1955; Stuart, 1963, 1969; Krishna, 1969; Howse, 1970; Roonwal, 1970; Watson & Gay, 1991; Thorne et al., 1993; Kambhampati & Eggleton, 2000), and agricultural (Harris, 1971; Lee & Wood, 1971; Scheffrahn & Su, 1992; Su & Scheffrahn, 2000) literature.

9. As *T. procerus* is the valid type species of both the genus *Termopsis* Heer, 1849 and the genus *Miotermes* Rosen, 1913, the name *Termopsis* (family TERMOPSIDAE) is formally a senior objective synonym of *Miotermes* (family MASTOTERMITIDAE).

10. Acceptance of this situation would be detrimental to the stability of termite nomenclature as all the species presently included in *Miotermes* would have to be transferred into the genus *Termopsis* and a new name would be needed for the genus that would include the species that had been included in *Termopsis*. The next available name is *Xestotermopsis* Rosen, 1913, a name that is currently considered to be a junior synonym of *Termopsis*. There would also be confusing complications associated with the family names TERMOPSIDAE and MASTOTERMITIDAE, as *Termopsis* is the type genus of TERMOPSIDAE. The name TERMOPSIDAE Holmgren, 1911 would become a junior synonym of MASTOTERMITIDAE Desneux, 1904 and a substitute name

would be needed for the family currently known as TERMOPSIDAE. The name STOLOTERMITINAE Holmgren, 1911 (p. 45) (currently used for a subfamily of TERMOPSIDAE) would be available, but its introduction at family rank would be confusing.

11. The International Commission on Zoological Nomenclature is accordingly asked:

- (1) to use its plenary power to set aside all previous fixations of type species for the nominal genus *Termopsis* Heer, 1849 and to designate *Termopsis breinii* Heer, 1849 as type species;
- (2) to place on the Official List of Generic Names in Zoology the following names:
 - (a) *Termopsis* Heer, 1849 (gender: masculine), type species by designation in (1) above *Termopsis breinii* Heer, 1849;
 - (b) *Mioterme*s Rosen, 1913 (gender: masculine), type species by original designation and monotypy *Termopsis procerus* Heer, 1849;
- (3) to place on the Official List of Specific Names in Zoology the following names:
 - (a) *breinii* Heer, 1849, as published in the binomen *Termopsis breinii* (specific name of the type species of *Termopsis* Heer, 1849);
 - (b) *procerus* Heer, 1849, as published in the binomen *Termopsis procerus* (specific name of the type species of *Mioterme*s Rosen, 1913);
- (4) to place on the Official List of Family-Group Names in Zoology the name TERMOPSIDAE Holmgren, 1911 (type genus *Termopsis* Heer, 1849).

Acknowledgements

We are grateful to V. Krishna and C.D. Michener for carefully reading earlier drafts of this petition and providing valuable commentary. This work was supported by a grant from the National Science Foundation, U.S.A. (DEB-9870097 to K. Krishna and D.A. Grimaldi).

References

- Banks, N. & Snyder, T.E. 1920. A revision of the Nearctic termites, with notes on biology and geographic distribution. *Bulletin of the United States National Museum*, **108**: 1–228.
- Cockerell, T.D.A. 1916. Insects in Burmese amber. *American Journal of Science*, **42**: 135–138.
- Emerson, A.E. 1933. A revision of the genera of fossil and recent Termopsinae (Isoptera). *University of California Publications in Entomology*, **6**: 165–195.
- Girard, M. 1879. *Les Insectes. Traité élémentaire d'entomologie* . . . , vol. 2. 1028 pp. Baillière, Paris.
- Hagen, H. 1854. Ueber die Neuropteren der Bernsteinfauna. *Verhandlungen des Zoologisch-botanischen Vereins in Wien*, **4**: 221–232.
- Hagen, H. 1858a. Monographie der Termiten. *Linnaea Entomologica*, **12**: 1–342.
- Hagen, H. 1858b. *Catalogue of the specimens of Neuropterous insects in the collection of the British Museum, Part I. Termitina*. 34 pp. British Museum, London.
- Handlirsch, A. 1907. *Die Fossilen Insekten und die Phylogenie der Rezenten Formen: Ein Handbuch für Paläontologen und Zoologen*. Pp. 641–1120. Engelmann, Leipzig.
- Harris, W.V. 1971. *Termites: their recognition and control*. xiii. 186 pp. Longmans, London.
- Heer, O. 1849. *Die Insektenfauna der Tertiärgebilde von Oeningen und von Radoboj in Croatien. Zweiter Theil: Heuschrecken, Flurfliegen, Aderflügler, Schmetterlinge und Fliegen*. 264 pp., 17 pls. Engelmann, Leipzig.
- Holmgren, N. 1911. Termitenstudien: 2. Systematik der Termiten. Die Familien Mastotermitidae, Protermitidae und Mesotermitidae. *Kungliga Svenska Vetenskaps-Akademiens Handlingar*, **46**(6): 1–88.

- Howse, P.E.** 1970. *Termites: a study in social behaviour*. 150 pp. Hutchinson University Library, London.
- Kambhampati, S. & Eggleton, P.** 2000. Taxonomy and phylogeny of termites. Pp. 1–23 in Abe, T., Bignell, D.E. & Higashi, M. (Eds.), *Termites: evolution, sociality, symbioses, ecology*. xxii, 466 pp. Kluwer Academic Publishers, Dordrecht.
- Krishna, K.** 1969. Introduction. Pp. 1–17 in Krishna, K. & Weesner, F.M. (Eds.), *Biology of termites*, vol. 1. xiii, 598 pp. Academic Press, New York.
- Krishna, K. & Grimaldi, D.** 2000. A new subfamily, genus, and species of termite (Isoptera) from New Jersey Cretaceous amber. Pp. 133–140 in Grimaldi, D. (Ed.), *Studies on Fossils in Amber, with Particular Reference to the Cretaceous of New Jersey*. viii, 498 pp. Backhuys, Leiden.
- Lee, K.E. & Wood, T.G.** 1971. *Termites and soils*. x, 251 pp. Academic Press, London.
- Nel, A. & Paicheler, J.-C.** 1993. Les Isoptera fossiles. État actuel connaissances, implications paléocéologiques et paléoclimatologiques [Insecta, Dictyoptera]. Pp. 103–179 in Nel, A., Martínez-Délclòs, X. & Paicheler, J.-C., *Essai de Révision des Aeschimoidea [Insecta, Odonata, Anisoptera] / Les Isoptera Fossiles [Insecta, Dictyoptera]*. 179 pp. CNRS Editions [Cahiers de Paléontologie], Paris.
- Pictet-Baraban, F.J. & Hagen, H.** 1856. Die im Bernstein befindlichen Neuropteren der Vorwelt. Pp. 41–126 in Berendt, G.C. (Ed.), *Die im Bernstein befindlichen organischen Reste der Vorwelt*, vol. 2. [2], 126 pp., 8 pls. Nicolaischen Buchhandlung, Berlin.
- Roonwal, M.L.** 1970. Termites of the Oriental region. Pp. 315–391 in Krishna, K. & Weesner, F.M. (Eds.), *Biology of Termites*, vol. 2. xiv, [1], 643 pp. Academic Press, London.
- Rosen, K. von.** 1913. Die fossilen Termiten: eine kurze Zusammenfassung der bis jetzt bekannten Funde. *Transactions of the Second International Congress of Entomology, Oxford 1912*, 2: 318–335.
- Scheffrahn, R.H. & Su, N.-Y.** 1992. Comparative toxicity of methyl bromide against ten Nearctic termite species (Isoptera: Termopsidae, Kalotermitidae, Rhinotermitidae). *Journal of Economic Entomology*, **85**: 845–847.
- Snyder, T.E.** 1949. Catalog of the termites (Isoptera) of the world. *Smithsonian Miscellaneous Collections*, **112**: 1–490.
- Stuart, A.M.** 1963. Studies on the communication of alarm in the termite *Zootermopsis nevadensis* (Hagen), Isoptera. *Physiological Zoology*, **36**: 85–96.
- Stuart, A.M.** 1969. Social behavior and communication. Pp. 193–232 in Krishna, K. & Weesner, F.M. (Eds.), *Biology of Termites*, vol. 1. xiii, 598 pp. Academic Press, New York.
- Su, N.-Y. & Scheffrahn, R.H.** 2000. Termites as pests of buildings. Pp. 437–453 in Abe, T., Bignell, D.E. & Higashi, M. (Eds.), *Termites: Evolution, Sociality, Symbioses, Ecology*. xxii, 466 pp. Kluwer Academic Publishers, Dordrecht.
- Thorne, B.L. & Carpenter, J.M.** 1992. Phylogeny of the Dictyoptera. *Systematic Entomology*, **17**: 253–268.
- Thorne, B.L., Grimaldi, D.A. & Krishna, K.** 2000. Early fossil history of the termites. Pp. 77–93 in Abe, T., Bignell, D.E. & Higashi, M. (Eds.), *Termites: evolution, sociality, symbioses, ecology*. xxii, 466 pp. Kluwer Academic Publishers, Dordrecht.
- Thorne, B.L., Haverty, M.L., Page, M. & Nutting, W.L.** 1993. Distribution and biogeography of the North American termite genus *Zootermopsis* (Isoptera: Termopsidae). *Annals of the Entomological Society of America*, **86**: 532–544.
- Wasmann, E.** 1897. Termiten von Madagaskar und Ostafrika. *Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft*, **21**: 137–182.
- Watson, J.A.L. & Gay, F.J.** 1991. Isoptera (termites). Pp. 330–347 in Naumann, I.D. (Ed.), *The insects of Australia: a textbook for students and research workers* [vol. 1]. xvi, [1], 542 pp. Cornell University Press, Ithaca.
- Weidner, H.** 1955. Körperbau, Systematik und Verbreitung der Termiten. Pp. 5–81 in Schmidt, H. (Ed.), *Die Termiten: Ihre Erkennungsmerkmale und wirtschaftliche Bedeutung*. 309 pp. Akademische Verlagsgesellschaft Geest und Portig, Leipzig.
- Weitschat, W. & Wichard, W.** 1998. *Atlas der Pflanzen und Tiere im Baltischen Bernstein*. 256 pp. Friedrich Pfeil, Munich.

- Weitschat, W. & Wichard, W. 2002. *Atlas of plants and animals in Baltic Amber*. 256 pp. Friedrich Pfeil, Munich.
- Wilson, E.O. 1971. *The insect societies*. x, [1], 548 pp. Harvard University Press, Cambridge, U.S.A.

Acknowledgement of receipt of this application was published in BZN 59: 161.

Comments on this case are invited for publication (subject to editing) in the *Bulletin*; they should be sent to the Executive Secretary, I.C.Z.N., c/o The Natural History Museum, Cromwell Road, London SW7 5BD, U.K. (e-mail: iczn@nhm.ac.uk).



Termopsis bremsii Heer, 1849 in Baltic Amber.